

Title Page

Title: Angiotensin II Receptors - Impact for COVID-19 Severity

Running Head: Adipose tissue and COVID-19

Category of the article: Review article

Number of words: 2672

Number of pages: 13 (including references)

Number of figures: 2

Number of tables: 0

Key Words: COVID-19, adipose tissue, obesity, ACE2, RAS

Authors: Aksoy H¹, Karadag AS¹, Wollina U²

Hasan Aksoy, Ayse Serap Karadag, Uwe Wollina

¹ Departments of Dermatology, Istanbul Medeniyet University, School of Medicine, Goztepe Training and Research Hospital, Istanbul, Turkey

² Department of Dermatology and Allergology, Städtisches Klinikum Dresden, Academic Teaching Hospital of the Technical University, Dresden, Germany

Correspondence:

Professor Uwe Wollina, Department of Dermatology and Allergology, Städtisches Klinikum Dresden
Academic, Teaching Hospital of the Technical University, Friedrichstrasse 41, 01067 Dresden, Germany
Email: Uwe.Wollina@klinikum-dresden.de.

ORCID

Ayşe Serap Karadag <https://orcid.org/0000-0003-4333-8274>

Uwe Wollina <https://orcid.org/0000-0001-5933-2913>

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/dth.13989

Abstract

COVID-19 is an outbreak of viral pneumonia which became a global health crisis, and the risk of morbidity and mortality of people with obesity are higher. SARS-CoV-2, the pathogen of COVID-19, enters into cells through binding to the Angiotensin Converting Enzyme (ACE) homolog-2 (ACE2). ACE2 is a regulator of two contrary pathways in renin angiotensin system (RAS): ACE-Ang-II-AT1R axis and ACE2-Ang 1-7-Mas axis. Viral entry process eventuate in downregulation of ACE2 and subsequent activation of ACE-Ang-II-AT1R axis. ACE-Ang II-AT1R axis increases lipid storage, reduces white-to-beige fat conversion and plays role in obesity. Conversely, adipose tissue is an important source of angiotensin, and obesity results in increased systemic RAS. ACE-Ang-II-AT1R axis, which has proinflammatory, profibrotic, prothrombotic and vasoconstrictive effects, is potential mechanism of more severe SARS-CoV-2 infection. The link between obesity and severe COVID-19 may be attributed to ACE2 consumption and subsequent ACE-Ang-II-AT1R axis activation. Therefore, patients with SARS-CoV-2 infection may benefit from therapeutic strategies that activate ACE2-Ang 1-7-Mas axis, such as Ang II reseptor blockers (ARBs), ACE inhibitors (ACEIs), Mas receptor agonists and ACE2.

Introduction

Coronavirus disease 2019 (COVID-19), is a pandemic resulted from the pathogen officially named severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2)¹. In line with World Health Organization (WHO), as of June 11, 2020, 7,273,958 cases and 413,372 deaths were confirmed worldwide². Obesity poses a higher risk for severe complications and mortality of SARS-CoV-2 infection for patients^{3,4}.

SARS-CoV-2 invades cells by attaching to the Angiotensin Converting Enzyme (ACE) homolog-2 (ACE2), a regulator of the renin angiotensin system (RAS)⁵. The consumption of ACE2 in the course of viral entry, subsequent increase in the angiotensin II and hyperactivation of the angiotensin II type 1 receptor (AT1R) appear to be major contributors to adverse outcomes in SARS-CoV-2 infection patients⁶⁻⁸.

Adipose tissue and RAS have a bidirectional relationship: Angiotensin II increases lipid storage and Ang II-AT1R axis is implicated in obesity, while adipose tissue is a crucial source of Ang II⁹⁻¹¹. Thus, the association between severe COVID-19 and obesity might be associated with destructive impact of Ang II hyperactivation and tendency to Ang II-AT1R axis. New therapeutic strategies may be generated in this context. Here we review complicated associations between adipose tissue, obesity, COVID-19 and RAS components, and we discuss potential treatment options for COVID-19.

The physiology of the angiotensin-2 receptors

The renin angiotensin system (RAS) is a systemic hormone system which acts in regulating blood pressure and energy homeostasis. RAS forms a complex cascade of pathways via which angiotensinogen (Agt) is sequentially cleaved by the enzymes renin, ACE and ACE2 into angiotensin I (Ang I), angiotensin II (Ang II), angiotensin 1–9 (Ang 1-9) and angiotensin 1-7 (Ang 1-7) (Figure 1)¹². Ang II, the crucial bioactive peptide of this system, mediates its cellular effects through two G-protein coupled receptors, Ang II Type 1 (AT1R) and Ang II Type 2 (AT2R) receptors, whereas effects of Ang 1-7 and Ang 1-9 are mediated by the Mas receptor. The AngII and Ang 1-7 peptides have mutual opposing effects. AngII-AT1R axis is proinflammatory, profibrotic, prothrombotic and has vasoconstrictive impacts, whereas Ang II-AT2R and Ang 1-7-Mas axes are anti-inflammatory, anti-fibrotic, anti-thrombotic and have vasodilatory impacts⁶.

Human adipocyte expresses Agt, renin, ACE, ACE2, AT1R, AT2R, and Mas receptor forming the local adipocyte RAS. Agt gene expression increases following feeding, long-chain fatty acids and hyperglycemia^{13,14}. Ang II raises lipogenesis in 3T3-L1 adipocytes via AT2R¹⁵. In addition to its lipogenic feature, Ang II prevents lipolysis in human white adipose tissue (WAT) in an AT1R-dependent way as well¹⁶⁻¹⁷. During the storage of excess energy, adipose tissue expansion appears by adipocyte hypertrophy (enlargement of adipocytes) and hyperplasia (adipogenesis). Adipocyte hypertrophy is linked with insulin resistance, whereas adipogenesis is a physiologically adaptive response to overnutrition and associated with better insulin-sensitivity¹⁸. Adipocyte-derived Agt acts actively in adipocyte differentiation. The Ang II-AT2R and Ang 1-7-Mas axes support adipocyte differentiation, while Ang II-AT1R axis applies anti-adipogenic impact in adipocytes from people with obesity¹⁹⁻²². Moreover, Ang II has negative impact on different steps of the insulin-signaling cascade and causes insulin resistance via AT1R. As a consequence, the net effect of Ang II and its receptors on adipose

tissue is to increase lipid storage and Ang II-AT1 axis is implicated in hypertension, obesity, and insulin resistance^{10,11}.

While the adipose tissue is affected by RAS, it also influences the systemic RAS. Thirty per cent of total circulating Agt is synthesized by the adipose tissue, being the second source of Agt after the liver⁹. Agt expression in WAT is improved in mice with obesity, resulting in increased systemic RAS²⁴. In a study, 44 healthy men that were overfed with diet enriched in lipid for 2 months demonstrated 1.5 and 1.4 fold increase in Agt gene expression and 1.2- and 1.5- fold increase in ACE expression in adipose tissue samples, after 2 and 8 weeks respectively²⁵. These data suggest that obesity influences adipose RAS and as consequence, components of systemic RAS.

SARS-CoV-2 entry into cells

The entry of SARS-CoV-2 into cells is mediated by the binding of the viral spike (S) protein to the ACE2, a regulator of the RAS^{5,26}. ACE2 is commonly bonded to cell membranes and merely exist in the circulation. Since the membrane-bounded ACE2 acts as a receptor for SARS-CoV-2 entry, in some publications it is called “ACE2 receptor”.

Firstly, N-terminal portion of the viral protein unit S1 binds to the ACE2 receptor, a trans-membrane type I glycoprotein. The second and crucial step for viral entry is the protein division between the S1 and S2 units, managed by the receptor transmembrane protease serine 2 (TMPRSS2)²⁷.

ACE2 receptors are explicit in the lungs, heart, vessels, gut, kidneys, testis, brain, and adipose tissue (epicardial adipocytes, adipocyte-like lung cells)^{28,29}. Also human keratinocytes express ACE2 and skin is a potential target for COVID-19, however, SARS-CoV-2 has not been detected in skin so far^{30,31}.

A disintegrin and metalloproteinase 17 (ADAM17), upregulated by Ang II via AT1R, mediates the cleavage of ACE2 from the cell membrane bound domain and release of ACE2 to the circulation³². SARS-CoV induces the downregulation of ACE2 membrane expression and plasma levels, through S protein induced ADAM17 activation and endocytosis of the ligand/receptor complex and subsequent to intracellular degradation of ACE2^{33,34}. Downregulation of ACE2 results in increased concentration of Ang II, which consecutively led to serious acute pulmonary injury, myocardial and endothelial dysfunction, obesity-associated hypertension, augmented inflammation, oxidative stress and coagulation⁶⁻⁸.

Obesity in COVID19 as a risk factor in general and for severe disease

It has been shown that obesity raises the vulnerability to infections; however, in cohort studies with SARS-CoV-2 disease, obesity prevalence was no higher than in general population^{3,4,35-37}. Since the

obesity is agreed to develop the risk of severe diseases, it can be assumed that obese individuals might be at risk of critical illness, if infected^{3,4,36}.

Obese COVID19 patients are more likely to be admitted to acute and critical care and have a risk for respiratory failure leading to invasive mechanical ventilation^{3,4}. In a retrospective analysis conducted on 112 cardiovascular disease patients infected by SARS-CoV-2, BMI was significantly higher in patients with a severe form of SARS-CoV-2 infection, compared with normal patients (25,5 [critical group] versus 22,0 [general group]; $P=0.003$). Moreover, among the non-survivor patient group, 88.2% of patients had a BMI $> 25 \text{ kg/m}^2$, which was a significantly higher proportion ($P < 0.001$) than in survivors (18.9%). Fulminant inflammation and thrombotic events were exacerbating reasons of death³⁸. In a study carried out in medical workers; compared to normal patients, BMI was significantly higher in patients with a severe form of SARS-CoV-2 infection (27.0 versus 22.0; $p<0.001$)³⁹. In another study with 5279 patients with laboratory confirmed SARS-Cov-2 infection, BMI $>40 \text{ kg/m}^2$ was one of the independent predictors for hospitalization⁴⁰. Among 383 patients from Shenzhen with SARS-CoV-2, the risk of developing severe infection was increased in overweight (BMI: $24.0\text{-}27.9 \text{ kg/m}^2$) and obese (BMI $\geq 28 \text{ kg/m}^2$) patients compared to normal weight (BMI: $18.5\text{-}23.9 \text{ kg/m}^2$) patients (overweight patients odds ratio [OR]:1.84, $p=0.05$; obese patients OR:3.40, $p=0.007$)⁴¹.

Obesity itself is a risk determinant for developing cardiac morbidity, and one of the supposed mechanisms is inflammation of epicardial adipose tissue (EAT). In mice on a high-fat diet, loss of ACE2 increases glucose intolerance and EAT inflammation, and administration of Ang 1-7 ameliorates EAT inflammation and cardiac steatosis, resulting in normalization of cardiac function⁴². In another study, obese female mice with adipocyte ACE2 deficiency, had an increased blood pressure response to Ang II⁴³. These studies suggest that ACE2-Ang 1-7-Mas axis, which is downregulated by SARS-CoV-2, has protective effects from obesity-induced cardiac dysfunction, glucose intolerance, hypertension and inflammation^{42,43}.

In terms of RNA level, the expression of ACE2 in adipose tissue was demonstrated to be higher than that in lung tissue, a major target tissue influenced by SARS-CoV-2⁴⁴. This result is surprising since it means adipose tissue can also be vulnerable to SARS-CoV-2 and can serve as a reservoir for the virus. Additionally, alveolar interstitium contains adipocyte-like cells called lipofibroblasts. Pulmonary lipofibroblasts can transdifferentiate into myofibroblasts, which may contribute to development of pulmonary fibrosis due to production of excessive extracellular matrix fibres and cause more severe COVID-19²⁹.

On the other hand obese individuals have more adipose tissue and thus have a higher number of ACE2-expressing and Ang II-producing cells, increased depletion of ACE2 during viral entry, subsequent larger amount of Ang II and unopposed functioning of ACE-Ang II-AT1 axis. Therefore, obesity might be a cause for increased adipose tissue derived circulating Agt, which is a potential culprit for a more severe SARS-CoV-2 infection (Figure 2).

In addition to impaired metabolic health, the criteria mediating the high risk of obese patients are believed to contain impaired respiratory mechanics, weakened immunity and pathogen defense, exuberant inflammation and chemotaxis which trigger “cytokine storm”, impaired fibrinolysis and subsequent thrombosis, vitamin D deficiency, and gut dysbiosis^{45,46}. In order to shed light to the relation between obesity and SARS-CoV-2 infection, further research is required.

Outlook for new treatment options

Since the SARS-CoV-2 infection causes downregulation of ACE2 and adipose tissue is an ample source of circulating Agt, the association between obesity and SARS-CoV-2 infection may be attributed to consumption of ACE2, and subsequent tendency to ACE-Ang II-AT1 axis. This hypothesis suggests potential therapeutic options for SARS-CoV-2 infection: Ang II receptor blockers (ARBs), ACE inhibitors (ACEIs), Mas receptor agonists, recombinant human ACE2 and ACE2 peptides as decoys for the virus.

ARBs/ACEIs treatment increases the expression of ACE2^{47,48}. Downregulation of ACE2 after SARS-CoV infection were shown to bring acute lung injury and beneficial net effects of ARB were displayed in SARS-CoV-induced acute lung injury in mice; thus we may suggest that treatment with ARBs/ACEIs or recombinant ACE2 would also protect from severe SARS-CoV-2 infection⁷.

A retrospective, multi-center research involving 1128 patients with hypertension and confirmed SARS-CoV-2 infection, revealed a lower mortality in the patients receiving ACEi/ARBs versus the other antihypertensive drugs (adjusted hazard ratio, 0.30 [95% CI, 0.12–0.70]; P=0.01). The difference remained significant following the adaptation for risk factors and baseline variables⁴⁹. In another study performed in 126 SARS-CoV-2 patients with hypertension, without statistical significance, lower proportion of critical patients (9.3% vs 22.9%; p=0.061) and a lower death rate (4.7% vs 13.3%; p=0.216), and significantly decreased concentrations of CRP (p=0.049) and procalcitonin (p=0.008) were monitored in ARBs/ACEIs group than non-ARBs/ACEIs group⁵⁰. Further randomized controlled trials (RCTs) are currently being conducted to better understand the influence of valsartan (NCT04335786), losartan (NCT04312009, NCT04311177, NCT04340557), intervention with various ARBs (NCT04394117), maintenance or discontinuation of previously used ARBs/ACEs treatment (NCT04364893, NCT04338009, NCT04351581) on morbidity and/or mortality in SARS-CoV-2 patients.

Recombinant ACE2, which has been approved as an experimental drug (APN01, GSK2586881) for adult respiratory distress syndrome (ARDS) and available for clinical trials, is another treatment option to replace ACE2. Animal models of acute lung failure^{51,52} have been used to show the efficacy of exogenous recombinant ACE2. Imai et al. demonstrated dramatically increased lung injury in ACE2 knockout mice, effects decreased by treating with recombinant human ACE2⁵¹. These findings extended to SARS-CoV-2, proposing for the use of ACE2 as a therapeutic agent in SARS-CoV-2, as well as providing soluble ACE2⁵³. Lately, SARS-CoV-2 was reported to directly infect engineered human blood vessel organoids and human kidney organoids, and recombinant human ACE2 marginally

Accepted Article

decreases the infectivity of SARS-CoV-2 in these ex-vivo models⁵⁴. ACE2 activates Ang 1-7-Mas axis and protects several organs by preventing fibrosis and lung injury as mentioned above, and it is likely that the circulating soluble form of ACE2 can also play a false receptor role for SARS-CoV-2, saturate the viral S-protein and may inhibit virus entry via membrane-bound ACE2.

ACE2 activators like diminazene aceturate, an antiparasitic drug mainly for veterinary use, or xanthenone can prevent pulmonary fibrosis, and also suggested as potential therapeutic agents for SARS-CoV-2 infection⁵⁵⁻⁵⁷.

TMPRSS2 is a membrane protease for ACE2 priming that is a vital step for the fusion of SARS-CoV-2 and target cell membranes and the resultant viral entry into the cells. The TMPRSS2 inhibitor camostat mesylate has been facilitated firstly to treat postoperative reflux esophagitis and for acute exacerbations of chronic pancreatitis, and has been reported to stop cellular infection by SARS-Cov-2⁵⁸. Currently, RCTs are being carried out to express whether camostat mesylate decreases SARS-COV-2 viral load in SARS-Cov-2 infection (NCT04353284, NCT04321096).

Activating the Ang 1-7-Mas axis may be another therapeutic approach to control the disease severity. Different Mas receptor agonists such as AVE 0991, hydroxypropyl β -cyclodextrin (HP β CD)-Ang 1-7, cyclic Ang 1-7, CGEN-856S and CGEN-857 have provided beneficial effects in hypertension, when tested in animal models⁵⁹. In a phase I/II study conducted with breast cancer patients, Ang 1-7 has been revealed to be safe, well tolerated and free of any dose-limiting toxicities⁶⁰. A randomized, placebo controlled clinical trial designed to determine the protective effect of TXA127 (a pharmacological formulation of Ang 1-7) on renal and pulmonary functions in patients with moderate to severe SARS-CoV-2 infection (NCT04401423).

Nevertheless, there are concerns whether therapeutic strategies based on ACE2 upregulation could be a threat for raised transmission of SARS-Cov-2^{53,61,62}. Some authors have suggested a biphasic effect of ACE2 upregulation, depending on clinical stage: detrimental in the contamination phase but beneficial in the tissue inflammation phase. However, ACE2 was shown to bind to SARS-CoV-2 with about 10- to 20-fold higher affinity than that bind to SARS-CoV and higher expression of ACE2 were not linked to higher susceptibility to infection with SARS-CoV^{63,64}. This may suggest that physiological expression of ACE2 may be already sufficient for SARS-CoV-2 infection, and further upregulation might not increase the risk.

Conclusion

Cellular entry process of SARS-CoV-2 downregulates ACE2, raises Ang II and enhances Ang II-AT1R axis, which has opposing effects to protective Ang 1-7-Mas axis. This is likely to be the cause of morbidity and mortality in patients with SARS-CoV-2 infection. Obesity, which is a risk factor for more severe COVID-19 disease, may contribute larger amounts of Ang II in the circulation and unopposed

functioning of ACE-Ang II-AT1 axis. There we may suggest that the link between adipose tissue/obesity and SARS-CoV-2 infection exists in the context of ACE2 depletion and RAS dysregulation. Therefore, Ang II receptor blockers (ARBs), ACE inhibitors (ACEIs), Mas receptor agonists and ACE2 seems to be potential therapeutic options for SARS-CoV-2 infection, peculiarly for individuals with obesity and metabolic syndrome.

References

1. Coronaviridae Study Group of the International Committee on Taxonomy of V. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. *Nat Microbiol.* 2020;5:536-544
2. WHO COVID-19 Dashboard. Available at: <https://covid19.who.int/>. Accessed 12 June 2020.
3. Lighter J, Phillips M, Hochman S, et al. Obesity in patients younger than 60 years is a risk factor for Covid-19 hospital admission [published online ahead of print, 2020 Apr 9]. *Clin Infect Dis.* 2020;ciaa415.
4. Goyal P, Choi JJ, Pinheiro LC, et al. Clinical Characteristics of Covid-19 in New York City [published online ahead of print, 2020 Apr 17]. *N Engl J Med.* 2020;NEJMc2010419.
5. Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature.* 2020;579:270-273.
6. Verdecchia P, Cavallini C, Spanevello A, Angeli F. The pivotal link between ACE2 deficiency and SARS-CoV-2 infection [published online ahead of print, 2020 Apr 20]. *Eur J Intern Med.* 2020;S0953-6205(20)30151-5.
7. Kuba K, Imai Y, Rao S, et al. A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury. *Nat Med.* 2005;11:875-879.
8. Kuba K, Imai Y, Penninger JM. Multiple functions of angiotensin-converting enzyme 2 and its relevance in cardiovascular diseases. *Circ J.* 2013;77:301-308.
9. Pahlavani M, Kalupahana NS, Ramalingam L, Moustaid-Moussa N. Regulation and Functions of the Renin-Angiotensin System in White and Brown Adipose Tissue. *Compr Physiol.* 2017;7:1137-1150. Published 2017 Sep 12.
10. Kalupahana NS, Massiera F, Quignard-Boulange A, et al. Overproduction of angiotensinogen from adipose tissue induces adipose inflammation, glucose intolerance, and insulin resistance. *Obesity (Silver Spring).* 2012;20:48-56.

11. Massiéra F, Bloch-Faure M, Ceiler D, et al. Adipose angiotensinogen is involved in adipose tissue growth and blood pressure regulation. *FASEB J*. 2001;15:2727-2729.
12. Chappell MC. Biochemical evaluation of the renin-angiotensin system: the good, bad, and absolute?. *Am J Physiol Heart Circ Physiol*. 2016;310:H137-H152.
13. Frederick RC Jr, Kahn BB, Peach MJ, Flier JS. Tissue-specific nutritional regulation of angiotensinogen in adipose tissue. *Hypertension*. 1992;19:339-344.
14. Safonova I, Aubert J, Negrel R, Ailhaud G. Regulation by fatty acids of angiotensinogen gene expression in preadipose cells. *Biochem J*. 1997;322:235-239.
15. Jones BH, Standridge MK, Moustaid N. Angiotensin II increases lipogenesis in 3T3-L1 and human adipose cells. *Endocrinology*. 1997;138:1512-1519.
16. Boschmann M, Ringel J, Klaus S, Sharma AM. Metabolic and hemodynamic response of adipose tissue to angiotensin II. *Obes Res*. 2001;9:486-491.
17. Goossens GH, Blaak EE, Arner P, Saris WH, van Baak MA. Angiotensin II: a hormone that affects lipid metabolism in adipose tissue. *Int J Obes (Lond)*. 2007;31:382-384.
18. Danforth E Jr. Failure of adipocyte differentiation causes type II diabetes mellitus?. *Nat Genet*. 2000;26:13.
19. Iwai M, Tomono Y, Inaba S, et al. AT2 receptor deficiency attenuates adipocyte differentiation and decreases adipocyte number in atherosclerotic mice. *Am J Hypertens*. 2009;22:784-791.
20. Shiuchi T, Cui TX, Wu L, et al. ACE inhibitor improves insulin resistance in diabetic mouse via bradykinin and NO. *Hypertension*. 2002;40:329-334.
21. Than A, Leow MK, Chen P. Control of adipogenesis by the autocrine interplays between angiotensin 1-7/Mas receptor and angiotensin II/AT1 receptor signaling pathways. *J Biol Chem*. 2013;288:15520-15531.
22. Brücher R, Cifuentes M, Acuña MJ, Albala C, Rojas CV. Larger anti-adipogenic effect of angiotensin II on omental preadipose cells of obese humans. *Obesity (Silver Spring)*. 2007;15:1643-1646.
23. Tsukuda K, Mogi M, Iwanami J, et al. Enhancement of Adipocyte Browning by Angiotensin II Type 1 Receptor Blockade. *PLoS One*. 2016;11:e0167704. Published 2016 Dec 19.
24. Yasue S, Masuzaki H, Okada S, et al. Adipose tissue-specific regulation of angiotensinogen in obese humans and mice: impact of nutritional status and adipocyte hypertrophy. *Am J Hypertens*. 2010;23:425-431.

25. Alligier M, Meugnier E, Debard C, et al. Subcutaneous adipose tissue remodeling during the initial phase of weight gain induced by overfeeding in humans. *J Clin Endocrinol Metab.* 2012;97:E183-E192.
26. Walls AC, Park YJ, Tortorici MA, Wall A, McGuire AT, Velesler D. Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. *Cell.* 2020;181:281-292.e6.
27. Glowacka I, Bertram S, Müller MA, et al. Evidence that TMPRSS2 activates the severe acute respiratory syndrome coronavirus spike protein for membrane fusion and reduces viral control by the humoral immune response. *J Virol.* 2011;85:4122-4134.
28. Gheblawi M, Wang K, Viveiros A, et al. Angiotensin-Converting Enzyme 2: SARS-CoV-2 Receptor and Regulator of the Renin-Angiotensin System: Celebrating the 20th Anniversary of the Discovery of ACE2. *Circ Res.* 2020;126:1456-1474.
29. Kruglikov IL, Scherer PE. The role of adipocytes and adipocyte-like cells in the severity of COVID-19 infections [published online ahead of print, 2020 Apr 27]. *Obesity (Silver Spring).* 2020;10.1002/oby.22856.
30. Xue X, Mi Z, Wang Z, Pang Z, Liu H, Zhang F. High expression of ACE2 on the keratinocytes reveals skin as a potential target for SARS-CoV-2 [published online ahead of print, 2020 May 23]. *J Invest Dermatol.* 2020;S0022-202X(20)31602-X.
31. Yao XH, Li TY, He ZC, et al. [A Pathological Report of Three COVID-19 Cases by Minimal Invasive Autopsies]. *Zhonghua Bing Li Xue Za Zhi.* 2020;49:411-417.
32. Patel VB, Clarke N, Wang Z, et al. Angiotensin II induced proteolytic cleavage of myocardial ACE2 is mediated by TACE/ADAM-17: a positive feedback mechanism in the RAS. *J Mol Cell Cardiol.* 2014;66:167-176.
33. Haga S, Yamamoto N, Nakai-Murakami C, et al. Modulation of TNF-alpha-converting enzyme by the spike protein of SARS-CoV and ACE2 induces TNF-alpha production and facilitates viral entry. *Proc Natl Acad Sci U S A.* 2008;105:7809-7814.
34. Wang S, Guo F, Liu K, et al. Endocytosis of the receptor-binding domain of SARS-CoV spike protein together with virus receptor ACE2. *Virus Res.* 2008;136:8-15.
35. Misumi I, Starmer J, Uchimura T, Beck MA, Magnuson T, Whitmire JK. Obesity Expands a Distinct Population of T Cells in Adipose Tissue and Increases Vulnerability to Infection. *Cell Rep.* 2019;27:514-524.e5.
36. Richardson S, Hirsch JS, Narasimhan M, et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area [published

online ahead of print, 2020 Apr 22] [published correction appears in doi: 10.1001/jama.2020.7681]. JAMA. 2020;323:2052-2059.

37. Buscemi S, Buscemi C, Batsis JA. There is a relationship between obesity and COVID-19 but more information is needed [published online ahead of print, 2020 May 12]. Obesity (Silver Spring). 2020;10.1002/oby.22883.

38. Peng YD, Meng K, Guan HQ, et al. Clinical characteristics and outcomes of 112 cardiovascular disease patients infected by 2019-nCoV. Zhonghua Xin Xue Guan Bing Za Zhi. 2020;48:E004.

39. Liu M, He P, Liu HG, et al. Clinical characteristics of 30 medical workers infected with new coronavirus pneumonia. Zhonghua Jie He He Hu Xi Za Zhi. 2020;43:E016.

40. Petrilli CM, Jones SA, Yang J, et al. Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: prospective cohort study. BMJ. 2020;369:m1966. Published 2020 May 22.

41. Cai Q, Chen F, Wang T, et al. Obesity and COVID-19 Severity in a Designated Hospital in Shenzhen, China [published online ahead of print, 2020 May 14]. Diabetes Care. 2020;dc200576.

42. Patel VB, Mori J, McLean BA, et al. ACE2 Deficiency Worsens Epicardial Adipose Tissue Inflammation and Cardiac Dysfunction in Response to Diet-Induced Obesity. Diabetes. 2016;65:85-95.

43. Shoemaker R, Tannock LR, Su W, et al. Adipocyte deficiency of ACE2 increases systolic blood pressures of obese female C57BL/6 mice. Biol Sex Differ. 2019;10:45.

44. Jia X, Yin C, Lu S, et al. Two Things about COVID-19 Might Need Attention. Preprints 2020, 2020020315 (doi: 10.20944/preprints202002.0315.v1).

45. Murugan AT, Sharma G. Obesity and respiratory diseases. Chron Respir Dis. 2008;5:233-242.

46. Muscogiuri G, Pugliese G, Barrea L, Savastano S, Colao A. Commentary: Obesity: The "Achilles heel" for COVID-19? [published online ahead of print, 2020 Apr 27]. Metabolism. 2020;108:154251.

47. Vuille-dit-Bille RN, Camargo SM, Emmenegger L, et al. Human intestine luminal Downloaded from <http://ahajournals.org> by on April 30, 2020 ACE2 and amino acid transporter expression increased by ACE-inhibitors. Amino Acids. 2015;47:693–705.

48. Li Y, Zeng Z, Li Y, et al. Angiotensin-converting enzyme inhibition attenuates lipopolysaccharide-induced lung injury by regulating the balance between angiotensin-converting enzyme and angiotensin-converting enzyme 2 and inhibiting mitogen-activated protein kinase activation. Shock. 2015;43:395–404.

49. Zhang P, Zhu L, Cai J, et al. Association of Inpatient Use of Angiotensin Converting Enzyme Inhibitors and Angiotensin II Receptor Blockers with Mortality Among Patients With Hypertension Hospitalized With COVID-19 [published online ahead of print, 2020 Apr 17]. *Circ Res*. 2020;10.1161/CIRCRESAHA.120.317134.
50. Yang G, Tan Z, Zhou L, et al. Effects Of ARBs And ACEIs On Virus Infection, Inflammatory Status And Clinical Outcomes In COVID-19 Patients With Hypertension: A Single Center Retrospective Study [published online ahead of print, 2020 Apr 29]. *Hypertension*. 2020;10.1161/HYPERTENSIONAHA.120.15143.
51. Imai Y, Kuba K, Rao S, et al. Angiotensin-converting enzyme 2 protects from severe acute lung failure. *Nature*. 2005;436:112-116.
52. Zou Z, Yan Y, Shu Y, et al. Angiotensin-converting enzyme 2 protects from lethal avian influenza A H5N1 infections. *Nat Commun*. 2014;5:3594. Published 2014 May 6.
53. Zhang H, Penninger JM, Li Y, Zhong N, Slutsky AS. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med*. 2020;46:586-590.
54. Monteil V, Kwon H, Prado P, et al. Inhibition of SARS-CoV-2 Infections in Engineered Human Tissues Using Clinical-Grade Soluble Human ACE2. *Cell*. 2020;181:905-913.e7.
55. Prata LO, Rodrigues CR, Martins JM, et al. Original Research: ACE2 activator associated with physical exercise potentiates the reduction of pulmonary fibrosis. *Exp Biol Med (Maywood)*. 2017;242:8-21.
56. da Silva Oliveira GL, de Freitas RM. Diminazene aceturate--An antiparasitic drug of antiquity: Advances in pharmacology & therapeutics. *Pharmacol Res*. 2015;102:138-157.
57. Rodríguez-Puertas R. ACE2 activators for the treatment of COVID 19 patients [published online ahead of print, 2020 May 7]. *J Med Virol*. 2020;10.1002/jmv.25992.
58. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell*. 2020;181:271-280.e8.
59. Povlsen AL, Grimm D, Wehland M, Infanger M, Krüger M. The Vasoactive Mas Receptor in Essential Hypertension. *J Clin Med*. 2020;9:267. Published 2020 Jan 18.
60. Rodgers KE, Oliver J, di Zerega GS. Phase I/II dose escalation study of angiotensin 1-7 [A(1-7)] administered before and after chemotherapy in patients with newly diagnosed breast cancer. *Cancer Chemother Pharmacol*. 2006;57:559-568.

63. Fang L, Karakiulakis G, Roth M. Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? [published correction appears in Lancet Respir Med. 2020 May 18;:]. Lancet Respir Med. 2020;8:e21.

64. Diaz JH. Hypothesis: angiotensin-converting enzyme inhibitors and angiotensin receptor blockers may increase the risk of severe COVID-19. J Travel Med. 2020;27:taaa041.

65. Dalan R, Bornstein SR, El-Armouche A, et al. The ACE-2 in COVID-19: Foe or Friend?. Horm Metab Res. 2020;52:257-263.

66. Perrotta F, Matera MG, Cazzola M, Bianco A. Severe respiratory SARS-CoV2 infection: Does ACE2 receptor matter? Respir Med. 2020;168:105996.

Figure Legends

Figure 1. Effects of Ang II and Ang 1-7 in the course of COVID-19.

Figure 2. Dysregulation of RAS in COVID-19 patients with obesity and potential link between obesity and more severe disease.




Fig. 1

42x32mm (762 x 762 DPI)




Fig. 2

42x32mm (762 x 762 DPI)